

THE INVASIVE CARDIOVASCULAR PROCEDURES AS A FACTOR FOR ECONOMIC EFFICIENCY IN HEALTHCARE

HRISTOVA Irinka¹
KOLEVA Greta¹
GEORGIEVA Despina¹

ABSTRACT

The healthcare establishes economical efficiency by reducing lethality of patients suffering from heart diseases. This is achieved with change in lifestyle, controlling the risk factors and the high chance of getting early diagnosed and treated with invasive cardiovascular procedures. Compared to 2010 mortality from cardiovascular diseases in Bulgaria has decreased by about 2%. Over time, invasive coronary procedures have increased in number and complexity. This trend is turning into an increased need for economic and human resources, among which the nursing staff is most affected. A number of factors can help to reduce lethality, including access to quality emergency care, timely patient transportation and high quality of service in specialized healthcare facilities for invasive diagnosis and treatment of cardiovascular disease. With the expansion of her role, the interventional radiological nurse should seek to continue her training and raise her level of competence. In Bulgaria, the doctors who work in invasive cardiovascular diagnostic laboratories, have acquired specialty of invasive cardiology according to Ordinance 1 of 2015 and in the same ordinance there is no specialty for nurses. They acquire the necessary practical skills and competences after starting their work in the nursing field. In the study of nurses training in Bulgaria, no specialized training in invasive diagnostics and treatment of cardiovascular diseases, is established. In the opinion of invasive cardiologist experts the procedural time and financial costs are decreased when working with a trained nurse in the team.

KEY WORDS *interventional radiological nurse, cardiovascular diseases, competencies, economic efficiency, healthcare.*

JEL: *I10, I19*

1. INTRODUCTION

Cardiovascular diseases (CVD), are the number one cause of death in the world: more people die annually from CVD, than for some other reason (WHO, 2016). A number of publications describe the burden of the cardiovascular diseases within Europe for 2016. Statistically they show that death from CVD, remains the most common cause of death on the old continent. It represents 45% of all deaths; 49% of women's deaths, and 40% among the men, as 1.4 million of these deaths are before the age of 75. Increasing hospitalization rates are linked to **rising costs** in healthcare systems that are connected with CVD, despite decreasing mortality. A number of factors can help reduce lethality, including access to quality emergency care, timely patient transportation and high quality care in specialized healthcare facilities for invasive diagnosis of cardiovascular disease (Townsend, N. et al., 2016).

¹ Asst. Prof., University of Ruse „Angel Kanchev“, Faculty of Public Health and Health Care, Department of Health Care, Bulgaria

According to data of the National Statistical Institute (NSI) of Bulgaria, for 2015, 72 028 Bulgarians died from diseases of the circulatory organs, which represents 65.4% of all causes of death in the country. Following the global trend, compared to 2010, the mortality rate from the CVD in our country decreased by about 2% (NSI, 2015). In the first six months of 2016, 5943 people in Bulgaria were diagnosed with acute myocardial infarction and 13.6% of them died (NCPHA, 2016).

In many European countries, cardiovascular mortality has continued to decrease over the last years. In high-income countries, the "turning point" is rapidly approaching when deaths from cancer will be more than the number of deaths from cardiovascular diseases, especially among men. The healthcare establishes economical efficiency by reducing lethality of patients suffering from heart diseases. This is achieved with change in lifestyle, controlling the risk factors and the high chance of getting early diagnosed and treated with invasive cardiovascular procedures.

2. STATEMENT

Over time, invasive coronary procedures have increased in number and complexity. This trend is turning into an increased need for economic and human resources, among which the nursing staff are most affected. In order to identify the possible predictors of nurses' workload, during and after diagnostic and interventional procedures, their commitment to serving patients in an Italian clinic has been investigated. A nurse in the Catheterization Laboratory has an average workload of 86 minutes and 174 minutes respectively for radial and femoral access. A nurse in a cardiac unit has an average workload of 386 minutes and 720 minutes respectively for radial and femoral access. The systematic application of radial access, when not contraindicated, is an effective strategy to reduce the workload of nurses during and after invasive coronary procedures (Amoroso, G. et al., 2005).

The advantages of the trans-radial approach compared to the trans-femoral are studied and scientifically proven. Radial access is increasingly preferred for both diagnostic and therapeutic procedures. The Allen test is used worldwide to determine whether the radial or ulnar artery passability is normal. This is done prior to the insertion of a radial cannula or catheterization of the radial or ulnar artery, as the presence of a catheter or other procedure-connected device in the artery, often results in thrombosis. A positive Allen test means that the patient does not have a double flush of the palm, so the Allen test is used to reduce the risk of hand ischemia (Physiopedia, 2015).

Trans-radial access is an increasingly standard approach, and the Allen test is needed. It is important that this practice varies between invasive diagnostic and therapy centers, and is usually not performed everywhere. One reason for this variability is that the sensitivity and specificity of the Allen test is relatively small (Valgimigli, M. et al., 2014). A variety of non-invasive options, including pulse oximetry and duplex ultrasound, are available to complement the Allen test, performed mainly by specially trained nurses. Yet, there is no consensus on the best test for circulatory hand safety, and the choice of test depends on the preference of the cardiologist and the available equipment. Given the high rate of false-positive Allen-test, many patients are wrongly excluded from trans-radial access for coronary angiography (Valgimigli, M. et al., 2015). In order to evaluate the efficacy and safety of the trans-radial approach, regardless of the results of the Allen test, a prospective data collection was conducted at the Cardiac Department at Red Cross Hospital, Athens, Greece in 2015. The study

concludes that the trans-radial approach for coronary angiography and angioplasty, can be effective and safe regardless of Alan's results before the procedure (Maniotis, C., 2015).

A large-scale randomized trial in England (London) proves the safety of patient's preparation for diagnostic cardiac catheterization, by specially trained nursing practitioners. The study compares the patient's preparation either by junior nurses or by nursing practitioners. Basic unwanted clinical events, patient's satisfaction, length of hospital stay, and cardiologists' assessment for patient preparation are been evaluated. This approach can be related to improving patient's satisfaction and reducing hospitalization time (Stables, R.H. et al., 2004).

Patient's satisfaction was also investigated in Iran, where a study was conducted among a group of patients indicated for coronary angiography. The qualitative study shows that patients have four main groups of expectations for coronary angiography: those from angiography, the treatment team, the need for angiography, education and training. Overall, the results of this study reveal a new look at the expectations of patients undergoing coronary angiography. Patient's education and training should include information addressed to expectations. Furthermore, an appropriate care program based on patient expectations, can promote quality of care and patient's satisfaction (Kalyani, M.N. et al., 2013).

Patients who have received preliminary information about the upcoming coronary angiography by the intervention-radiologist nurse, and have slept well before the procedure show an extremely low level of discomfort. Pre-information and patient's training should be provided by experienced nursing practitioners (Park, J. et al., 2009).

Coronary angiography is a widely used invasive procedure for the diagnosis and therapy of coronary artery disease. Preliminary information and patient's education is essential. The results of the study aimed at assessing the effectiveness of two educational methods (video information and oral information) for patients with imminent coronary angiography, confirm the usefulness of video information prior to invasive coronary angiography. The group of patients who also received video information, showed statistically significant reductions in heart rate and blood pressure after educational activity, and had significantly higher levels of comfort, satisfaction and tolerance, than the control group receiving only routine verbal information from the nurses (Jamshidi, N. et al., 2013)

Patients who wait for coronary angiography are often worried and anxious. They experience significant negative emotions before the procedure, which may lead to an increase in blood pressure, heart rate and respiratory rate. Such non-adaptive responses can not only increase patients' need for sedative drugs, but also increase the length of post-angiographic hospitalization. Therefore, it is important to implement some supportive actions, to reduce patient's anxiety and stabilize their vital signs before coronary angiography. In a conducted study to investigate the effect of the multimodal preparation package for patients for imminent coronary angiography, heart rate, systolic and diastolic blood pressure values, were evaluated. The study showed that multimodal packet preparation is effective in lowering blood pressure and heart rate. It is therefore advisable to use a combined complex preparation of the patient, with imminent coronary angiography (Adib Hajbaghery, M. et al., 2014).

A Swiss study, designed to assess the learning outcomes and those of the patient's percutaneous coronary intervention (PCI) trial, shows that current preventative practice almost fails to meet the challenge posed by advances in modern invasive

cardiology. The results of the post-PCI study of the patients motivate a deeper review and adaptation of cardiac rehabilitation programs, with the inclusion of trained nurses to improve the patient's understanding of the disease and cardio-protective lifestyle (Perk, A.R. et al., 2015).

Various educational interventions are being implemented to help patients with cardiovascular disease, maintain their health, develop self-care skills, and reduce the number and duration of re-hospitalizations. The effectiveness of multimedia intervention, combined with nursing intervention, in patients with heart failure has been studied in an Australian hospital. The in-patient education of these patients has been positive, and approaches to patients include: needs assessment and application of an individualized plan. Multimedia resources and learning strategies are effective in improving patients' therapeutic outcomes (Boyde M., 2015)

In the "Guidelines for Nursing Interventional Radiology", it is stated that the preliminary assessment of the admittance of planned patients, is an established role of the interventional radiological nurse. Nurses must have a different level of qualification, corresponding to the assumed responsibility, and this must be noted in their job description. Interventional nurses must have basic knowledge of resuscitation and continue their training in resuscitation techniques. Nursing tasks do not release the clinical radiologist from overall responsibility to ensure that all necessary actions are taken. The primary responsibility of the nurse in interventional radiology, is to observe the patient and ensure his safety (Guidelines for Nursing Care in Interventional Radiology, 2014).

Coordinated protocols and integrated means of treatment should be established, so that the interventional radiological nurse can judge whether the patient is suitable for ambulatory, or stationary interventional treatment. This will help to avoid unnecessary and sometimes prolonged hospitalizations, and thus optimize the resources and potential delay of the patient's treatment. During the preliminary assessment, the nurse prepares routine research and, according to the results, acts on accordant protocols. The nurse must have knowledge of all intervention procedures, so that she can inform the patient, and be able to answer all his questions.

Within the intervention itself, the role of the interventional radiologist nurse is to support the procedure and, with the other members of the multidisciplinary team, to take care of the patient during the study. Two experienced nurses are required to perform complex procedures - one to work sterilely with the clinical radiologist during the procedure, and the other to observe the patient and perform the appointments of the physician. The second nurse should also provide the necessary devices during the procedure. Interventional nurses should have appropriate competencies for cannulation of vein and intravenous opiate administration. With the expansion of her role, the interventional radiological nurse should seek to continue her training, and raise her level of competence. Specialized nurses in radiology are well acquainted with the special equipment and medical supplies, and can provide the necessary devices for each procedure. Early recognition of allergic reactions to the contrast matter is also part of the nurse's competence. The interventional radiological nurse evaluates, plans, and implements general nursing care for the patient (including pulse monitoring, blood pressure, electrocardiogram, oxygen saturation of the blood, etc.). She co-ordinates and documents the planned patient care, during the invasive procedure. The nurse creates a safe and therapeutic environment for both the patient and the staff. She also complies with the requirements for safe work in X-ray environment, practice aseptic principles

and acting as a leading professional in the control of infections. Communicates effectively with other members of the Interventional radiology team.

The unique and patients following the procedure require high qualification and specialization of intervention nurses.

According to the Guidelines for Nursing in Interventional Radiology, developed by the Royal College of Nursing and the Royal College of Radiology, a minimum ratio of staff comprising two nurses - one them experienced, with appropriate qualification.

Invasive cardiology in Bulgaria made its first steps in 1963, when a functional sector of Blood Diagnostics, was formed at the National Cardiology Hospital. In 1975, the first coronarography in Bulgaria was made, which is A. Savova's merit (Tomov, I., 2002). The first invasive manipulation, in the sense of a healing procedure in Bulgaria, was carried out in 1986 by Prof. Arnaudov and Assis. Tsonzarova. This is a child's pulmonary valve dilation. While in adults the first interventions were performed about two years later - coronary valvular dilations (Naydenova, I., 2015). The name of Prof. Bozhidar Finkov should be written in the history of Bulgarian medicine, because in 1991, he performed the first cardio-invasive manipulation in our country. He put the first stent in Bulgaria, on a patient with anterior heart attack and saved him. Later on, he performed the first valvuloplasty in Bulgaria (UMHAT St. Anna - Sofia AD, 2016).

In 2002, Dr. A. Doganov introduces the more gentle and economical trans-radial access. Trans-radial access quickly becomes routine, and is used for both diagnostic and interventional procedures, including patients with acute myocardial infarction (Tomov, I., 2002). Trans-radial access for cardiac catheterization, which emerged as an alternative to the trans-femoral, is now a "gold standard" for many catheterization laboratories (Krastev, V., 2005).

When an independent analysis of the data of the National Health Insurance Fund was carried out in 2008, the independent consultancy company Sanigest found that percutaneous coronary angioplasty in Bulgaria, is nearly twice as low as in the developed European countries, which objectively indicates the potential problems - with the access to specialized health care, with diagnostics of patients and the need for such a procedure, as well as the shortage of hospital structures for invasive cardiovascular diagnostics (Vekov, T., 2009).

For just five years, from the beginning of 2010 to 2014, in Bulgaria, the number of hospitals funded by the National Health Insurance Fund on clinical pathways for invasive cardiology, increased from 33 to 50 (Bozukova, M., 2015). By the beginning of 2016, the clinics of Invasive Cardiology in Bulgaria are 71 (Panev, D, 2016), 10 of which (MH, 2016) function in Sofia only.

For a short period of time, the wards for invasive cardiovascular diagnostics have increased many times, requiring medical teams to be redirected from cardiac structures, and recruited to work in catheterization laboratories for non-invasive diagnostics and therapy. The rapid pace of expansion of high-tech clinics for invasive cardiology in practice, requires an adequate response from universities to the training of medical staff.

3. CONCLUSIONS

Invasive percutaneous procedures are proven to be financially more effective compared to known surgical methods for the treatment of cardiovascular diseases. For patients with cardiogenic shock who need hemodynamic support, percutaneous resources of supporting the cardiovascular system may be used at an earlier stage, which

contributes better results, shorter hospital stay and lower financial costs for the treatment compared to traditional alternatives for surgical intervention.

Recent years in invasive cardiovascular procedures, it is increasingly preferred trans-radial access for cardiac catheterization but, special training is needed for the medical and nursing staff. In recent years, it has become increasingly important as a first choice due to the minimal possible complications for the patient and the drastic reduction in financial costs for medical consumables and hospital stay.

In Bulgaria, the doctors working in the invasive cardiovascular diagnostic laboratories have acquired specialty of invasive cardiology according to Ordinance 1 and no specialty for nurses is prescribed in the same ordinance (Naredba No. 1, 2015). They acquire the necessary practical skills and competencies after starting their work in the nursing field. In the research of nurses training in Bulgaria, no specialized training in invasive diagnostics and treatment of cardiovascular diseases, is found. The Bulgarian Association of Healthcare Professionals organizes, coordinates, conducts and records the continuing training of nurses, midwives and associated medical specialists, according to Art. 182 of the Health Act. In the plan for the continuing training of nurses for 2017, only two of the universities in Bulgaria offer training, related to invasive cardiology – Angel Kanchev University of Ruse and Medical University of Sofia, which is not directed to access to the procedure - trans-radial or trans-femoral. However, in all hospitals for invasive cardiology in Bulgaria the trans-radial access is the first choice when it is possible and there is no contraindication for it. Led by financial profitability and minimal possible complications for patients. Leading invasive cardiologists in our country prefer it. According to data from the Specialized Hospital for Active Cardiology Treatment in Rousse, Bulgaria invasive procedures through trans-radial access are 93.1% of all procedures for 2007 which contributes to financial stability and competitiveness.

Financial costs in modern healthcare are one of the "turning points" in managing treatment centers. In the opinion of invasive cardiologist experts procedural time and financial costs are increased when working with an untrained nurse (Hristova. I., 2016).

The specific opportunities offered by invasive cardiology to reduce the cost of treating patients with cardiovascular disease are the subject of further research, opportunities are being put in place to introduce new innovative practices into the healthcare system. The development of cost-effective treatment approaches is a priority for every medical institution.

REFERENCES

- Adib Hajbaghery, M., Moradi, T., Mohseni R. (2014) "Effects of a multimodal preparation package on vital signs of patients waiting for coronary angiography". Available on-line at <http://www.ncbi.nlm.nih.gov/pubmed/25414893>
- Amoroso, G. et al. (2005) "Clinical and procedural predictors of nurse workload during and after invasive coronary procedures: the potential benefit of a systematic radial access". *European Journal of Cardiovascular Nursing* 4 (2005) 234 – 241.
- Boyde, M., Peters, R., Hwang, R. et al. (2015) "Intervention for Patients With Heart Failure: A Study Protocol". Available on-line at <http://www.ncbi.nlm.nih.gov/pubmed/26696034>
- Bozukova, M. (2015) NZOK "Spira krancheto" za invazivnite izsledvania na zdravi patsienti". <http://www.mediapool.bg/nzok-spira-krancheto-za-invazivnite-izsledvaniya-na-zdravi-patsienti-news230656.html>
- Hristova, Ir., (2015) "Otsenka na arterialnoto kravosnabdyavane na dlanta chrez test na Allen",

- Studentska nauchna konferentsia na RU 2015, Nauchni trudove, 2015, стр. 132, ISSN 1311-3321
- Jamshidi, N., Abbaszadeh, A., Kalyani, M.N. et al. (2013) "Effectiveness of video information on coronary angiography patients' outcomes". Available on-line at <http://www.ncbi.nlm.nih.gov/pubmed/24151693>
- Kalyani, M.N., Sharif, F., Ahmadi, F. et al. (2013) "Iranian patient's expectations about coronary angiography: A qualitative study.", Available on-line at <http://www.ncbi.nlm.nih.gov/pubmed/23983751>
- Krastev, V. (2005) "Transradialna koronarna angiografija i angioplastika, Profilaktika diagnostika terapija aktualni problemi". Sofia, ISBN 954 – 91788 – 1. 419-423.
- Maniotis, C., Koutouzis, M., Andreou, C. et al. (2015) "Transradial Approach for Cardiac Catheterization in Patients With Negative Allen's Test. J Invasive Cardiol". 27(9):416-20. Available on-line at <https://www.ncbi.nlm.nih.gov/pubmed/?term=Transradial+Approach+for+Cardiac+Catheterization+in+Patients+With+Negative+Allen%E2%80%99s+Test>
- Ministerstvo na zdraveopazvaneto, (2016) "Lechebni zavedenia za bolnichna pomosht, dispanseri", DMSGD, TsSMP, TsTH. Available on-line at <http://www.mh.government.bg/bg/administrativni-uslugi/registri/lechebni-zavedeniya/>
- Naredba №1 ot 22.01.2015 g. za pridobivane na spetsialnost v sistemata na zdraveopazvaneto. Available on-line at http://www.mu-pleven.bg/forms/Naredba_1_SDO.pdf
- Natsionalen Statisticheski Institut: Naselenieto na Bulgaria prodalzhava da namalyava i zastaryava. Available on-line at <http://www.bgues.com/bylgaria/obshchestvo/4423760/>
- Natsionalen Statisticheski Institut (2015) Umirania po prichini za smartta prez 2015 g. po pol i vazrastovi grupi. Available on-line at <http://www.nsi.bg/bg/content/3351>
- Natsionalen tsentar po obshtestveno zdrave i analizi (2015) Godishna informatsia. Available on-line at http://ncphp.government.bg/index.php?option=com_content&view=category&layout=blog&id=96&Itemid=640&lang=bg
- Naydenova, I. (2015) Dots. Bozhidar Finkov: Invazivnata kardiologia e dobre razvita u nas. Available on-line at <http://www.bestdoctors.bg/article/id/1056406>
- Panev, D. (2016) Zakrivat 48 ot 71 Kliniki po invazivna kardiologia. Available on-line at http://dariknews.bg/view_article.php?article_id=1549754
- Park, J. Choi. (2009) "Factors predicting patient discomfort after coronary angiography". Available on-line at <http://www.ncbi.nlm.nih.gov/pubmed/20071899>
- Perk, A.R., Hambraeus, K., Burell G. et al. (2015) "Study of Patient Information after percutaneous Coronary Intervention (SPICI): should prevention programmes become more effective". Available on-line at <http://www.ncbi.nlm.nih.gov/pubmed/24472705>
- Physiopedia. Allen test. Available on-line at http://www.physio-pedia.com/Allen_Test
- Stables, R.H., Booth, J., Welstand, J. (2004) "A randomised controlled trial to compare a nurse practitioner to medical staff in the preparation of patients for diagnostic cardiac catheterisation: the study of nursing intervention in practice (SNIP)". Available on-line at <http://www.ncbi.nlm.nih.gov/pubmed/15053888>
- The Royal College of Nursing and The Royal College of Radiologists. (2014) "Guidelines for nursing care in interventional radiology, Second edition". London, Ref No. BFCR(14)7, https://www.rcr.ac.uk/system/files/publication/field_publication_files/BFCR%2814%297_Guidelines_nursing_interventional_radiology.pdf
- Tomov, I. (2002) Kardiologia, Parvo izdanie. Znanie. 667-714.
- Townsend, N., Wilson, L., Bhatnagar, P., et al. (2016) "Cardiovascular disease in Europe: epidemiological update" 2016. Available on-line at <http://eurheartj.oxfordjournals.org/content/early/2016/08/10/eurheartj.ewh334>
- UMBAL Sveta Anna – Sofia AD. Klinika po kardiologia. Available on-line at <http://www.sveta-anna.eu/kardiologia.html>

- Valgimigli, M., Campo, G., Penzo, C. et al. (2014) "Transradial coronary catheterization and intervention across the whole spectrum of Allen test results". *J Am Coll Cardiol.* 63:1833–41. Available on-line at <https://www.ncbi.nlm.nih.gov/pubmed/24583305>.
- Valgimigli, M., Gagnor, A., Calabro, P. et al. (2015) "Radial versus femoral access in patients with acute coronary syndromes undergoing invasive management: a randomized multicentre trial". *Lancet.* 385:2465–76.
- Vekov, T. (2009) "Mezhdunarodniyat i balgarskiyat opit v oblastta na invazivnatai intervensionalna kardiologia – razvitie, neobhodimost, rezultati". MU – Sofia, Sardechno-sadovi zabolyavania. 40, 2009.
- WHO. (2016) "Cardiovascular diseases". Available on-line at <http://www.who.int/mediacentre/factsheets/fs317/en/>